# **JTGGA CME/CPD CREDITING**



## Answer form for the article titled "*The possible role of the da Vinci robot for patients with vulval carcinoma undergoing inguinal lymph node dissection*" within the scope of CME/CPD

#### 1. Which of the following is considered the gold standard of vulval cancer management?

- a) Radical vulvectomy or wide local excision plus inguinal lymphadenectomy (triple incision technique).
- b) Radical vulvectomy or wide local excision plus inguinal lymphadenectomy and pelvic lymphadenectomy.
- c) Radical vulvectomy or wide local excision plus inguinal lymphadenectomy (butterfly technique).
- d) Radical vulvectomy or wide local excision followed by sentinel lymph node technique
- e) Radical vulvectomy or wide local excision followed by robotic lymphadenectomy

#### 2. Which of the following complications is not related to inguinal lymphadenectomy?

- a) Lymphocyst
- b) Lymphedema
- c) Skin flap necrosis
- d) Urinary incontinence
- e) Wound infection

#### 3. Which of the following is not an advantage of robotic approach?

- a) Comfort for the surgeon
- b) Shorter surgical time
- c) 3D approach
- d) High magnification
- e) Instruments with a higher degree of freedom

#### 4. Which of the following is false regarding the role of minimal invasive approach for inguinal lymphadenectomy?

- a) Shorter hospital stay
- b) Shorter recovery period
- c) Equally safe and oncologically effective
- d) Lower morbidity rates
- e) Less postoperative pain

#### 5. Which of the following is not true regarding the steps of robotic inguinal lymphadenectomy?

- a) Femoral triangle is identified and a 2-cm incision is performed about 3 cm below its inferior aspect
- b) Scarpa's fascia is identified and after blunt-finger dissection, the scope is used to create a superficial subcutaneous flap by sweeping the lens under the fascia
- c) Pneumoperitoneum up to 5-10 mm Hg pressure
- d) Bipolar Maryland and monopolar scissors are the main instruments used
- e) Saphenofemoral junction is exposed after opening the fascia lata and deep pelvic lymph node dissection can also be performed if necessary

#### 6. Which of the following is not correct?

- a) Prospective randomized controlled trials are necessary to clarify the morbidity rates and advantages of robotic inguinal lymphadenectomy for vulval cancer patients
- b) The main disadvantage is the high cost
- c) Objections could be raised regarding the learning curve
- d) The risk of port site metastasis is not clear
- e) Robotic approach at the moment is cost effective for the health systems

# JTGGA CME/CPD CREDITING



Questions on the article titled "*The possible role of the da Vinci* robot for patients with vulval carcinoma undergoing inguinal lymph node dissection" within the scope of CME/CPD

1 <sup>st</sup> Question						4th Question					
A	В	С	D	Е		A	В	С	D	Е	
2 <sup>nd</sup> Question						5 <sup>th</sup> Question					
A	В	С	D	Е		A	В	С	D	Е	
3rd Ques	tion					6 <sup>th</sup> Question					
A	В	С	D	E		Α	В	С	D	E	

People who answer these questions will receive "2 TMA-CME/CPD credits"

### TMA-CME CREDITING BOARD ENQUIRY FORM

### JTGGA MANUSCRIPT 2017/2

TR Identification Number	This form will not be reviewed if TR Identification Number is not stated.
Name	
Surname	
Signature	The City You Work In
Your Institution	

**IMPORTANT NOTE:** You may apply for Turkish Medical Association CME/CPD credits by answering the questions in the front page, filling in your personal information and sending this form to "Abdi İpekçi Cad. No: 2/7 34367 Nişantaşı, İstanbul" by post. This form should arrive to the above-mentioned address latest by August 31, 2017.